

AN EXTENSION OF WICK'S THEOREM

C. VIGNAT AND S. BHATNAGAR

ABSTRACT. We propose an extension of a result by Repetowicz et al. [1] about Wick's theorem and its applications: we first show that Wick's theorem can be extended to the uniform distribution on the sphere and then to the whole class of elliptical distributions. Then, as a special case, we detail this result for distributions that are Gaussian scale mixtures. Finally, we show that these results allow to recover easily a theorem by Folland [5] about integration of polynomials over the sphere.

1. WICK'S THEOREM

In quantum field theory [2], the determination of the partition function of some systems involves the computation of integrals of the form

$$\int_{\mathbb{R}^n} P(X) \exp\left(-\frac{X^t \Sigma^{-1} X}{2}\right) dX,$$

where $P(X)$ is a monomial in the components of $X \in \mathbb{R}^n$, that is, in a probabilistic approach, mixed moments of a Gaussian random vector with covariance matrix Σ .

Wick's theorem provides a simple formula for these moments: let us denote by X a Gaussian vector in $\mathbb{R}^n$ with zero mean and covariance matrix Σ ; a linear form $\mathcal{L}_i$ on $\mathbb{R}^n$ is identified with the vector of its coefficients

$$\mathcal{L}_i(X) = l_i^t X$$

A *pairing* σ on the set $I_{2m} = \{1, \dots, 2m\}$ is a partition of I_{2m} into m disjoint pairs; the set Π_{2m} of pairings of I_{2m} is composed of $\frac{2m!}{2^m m!}$ elements. For example, in the case $m = 2$, there are 3 different pairings on $I_4 = \{1, 2, 3, 4\}$, namely $\sigma_1 = \{(1, 2), (3, 4)\}$, $\sigma_2 = \{(1, 3), (2, 4)\}$ and $\sigma_3 = \{(1, 4), (2, 3)\}$. We denote by I_{2m}/σ the set of indices i such that $\sigma = \{(i, \sigma(i))\}$.

Wick's theorem states as follows [2].

Theorem. *Let m be an integer then*

$$E \prod_{i=1}^{2m} \mathcal{L}_i(X) = \sum_{\sigma \in \Pi_{2m}} \prod_{i \in I_{2m}/\sigma} l_i^t \Sigma l_{\sigma(i)}$$

If $2m$ is replaced by $2m + 1$, then the expectation above equals zero.

As an example, with $m = 2$ and $\mathcal{L}_i(X) = X_i$, we obtain

$$E[X_1 X_2 X_3 X_4] = E[X_1 X_2] E[X_3 X_4] + E[X_1 X_3] E[X_2 X_4] + E[X_1 X_4] E[X_2 X_3].$$

2. SPHERICAL AND ELLIPTICAL DISTRIBUTIONS

In this paper, we extend Wick's theorem to the class of spherical random vectors: such a vector $X \in \mathbb{R}^n$ is characterized [3, Ch.2] by the fact that its characteristic function $\psi(Z) = E \exp(iZ^t X)$ is invariant by orthogonal transformation; in other words,

$$\psi(Z) = \phi(Z^t Z)$$

for some function $\phi : \mathbb{R}^+ \rightarrow \mathbb{C}$. In this case, a stochastic representation of X writes

$$X = rU$$

where U is a random vector uniformly distributed on the unit sphere $\mathcal{S}_n = \{X \in \mathbb{R}^n; \|X\| = 1\}$ and $r \in \mathbb{R}^+$ is a positive random variable independent of U . If moreover vector X admits a density f_X , then it is necessarily of the form

$$f_X(X) = g(X^t X)$$

for some function $g : \mathbb{R} \rightarrow \mathbb{R}^+$.

An extension of the spherical property is the elliptical property [3, Ch.2]: a random vector $Y \in \mathbb{R}^n$ is elliptical with $(n \times n)$ characteristic matrix Σ if

$$(1) \quad Y = \Sigma^{1/2} X$$

where X is spherical. We assume without loss of generality that all vectors have zero mean.

3. WICK'S THEOREM FOR THE UNIFORM DISTRIBUTION ON THE SPHERE

Wick's theorem can be extended to the case of a vector uniformly distributed on the sphere $\mathcal{S}_n$ as follows.

Theorem. *If U is uniformly distributed on the sphere $\mathcal{S}_n$ then*

$$E \prod_{i=1}^{2m} \mathcal{L}_i(U) = \frac{\Gamma(\frac{n}{2})}{2^m \Gamma(m + \frac{n}{2})} \sum_{\sigma} \prod_{i \in I_{2m}/\sigma} l_i^t l_{\sigma(i)}$$

Proof. A stochastic representation for vector U is

$$U = \frac{N}{\|N\|}$$

where N is a Gaussian vector with unit covariance matrix. By the polar factorization property, random variable $r = \|N\|$ is independent of U ; we deduce from the linearity of $\mathcal{L}_i$ that

$$E \prod_{i=1}^{2m} \mathcal{L}_i(N) = E \prod_{i=1}^{2m} r \mathcal{L}_i(U) = E r^{2m} E \prod_{i=1}^{2m} \mathcal{L}_i(U)$$

so that

$$E \prod_{i=1}^{2m} \mathcal{L}_i(U) = \frac{E \prod_{i=1}^{2m} \mathcal{L}_i(N)}{E r^{2m}}.$$

Since $r = \sqrt{N^t N}$ is chi-distributed with n degrees of freedom, elementary algebra yields

$$Er^{2m} = \frac{2^m \Gamma(m + \frac{n}{2})}{\Gamma(\frac{n}{2})}.$$

□

From the linearity of forms $\mathcal{L}_i$ and by stochastic representation (1), we deduce a version of Wick's theorem for any elliptical vector as follows.

Theorem. *If $X \in \mathbb{R}^n$ is an elliptical vector with stochastic representation*

$$(2) \quad X = a\Sigma^{1/2}U$$

and if random variable a has moments up to order $2m$, then

$$E \prod_{i=1}^{2m} \mathcal{L}_i(X) = \frac{Ea^{2m}}{Er^{2m}} \sum_{\sigma} \prod_{i \in I_{2m}/\sigma} l_i^t \Sigma l_{\sigma(i)}$$

4. WICK'S THEOREM FOR GAUSSIAN SCALE MIXTURES

A random vector X is a Gaussian scale mixture if its probability distribution writes

$$f_X(X) = \int_0^{+\infty} g(x; a\Sigma) dH(a)$$

where $g(x; \Sigma)$ is the Gaussian distribution with $(n \times n)$ covariance matrix Σ on $\mathbb{R}^n$ and H is a cumulative distribution function. A stochastic representation of such a vector is

$$(3) \quad X = \frac{N}{\sqrt{a}}$$

where N is a Gaussian vector with covariance matrix Σ and a is a scalar random variable independent of N with cumulative distribution function H . Cauchy distributions, alpha-stable and Student-t distributions belong to the family of Gaussian scale mixture distributions.

If X is a Gaussian scale mixture, then it is also elliptical; however, the contrary is not true, as shown [4] by the Pearson II (or Student-r) family of distributions

$$f_X(X) = \frac{\Gamma\left(\frac{2-q}{1-q} + \frac{n}{2}\right)}{\Gamma\left(\frac{2-q}{1-q}\right) |\pi p K|^{1/2}} \left(1 - \frac{X^t K^{-1} X}{p}\right)_+^{\frac{1}{1-q}}$$

with $p = n + 2\frac{2-q}{1-q} > 0$ and notation $(x)_+ = \max(x, 0)$.

Using stochastic representation (3), Wick's theorem extends to Gaussian scale mixtures as follows.

Theorem. *If X is a Gaussian scale mixture as in (3) with $Ea^{-m} < \infty$, then*

$$E \prod_{i=1}^{2m} \mathcal{L}_i(X) = E[a^{-m}] \sum_{\sigma \in \Pi_m} \prod_{i \in I_{2m}/\sigma} l_i^t \Sigma l_{\sigma(i)}$$

We note that Repetowitz et al. [1] deal only with the case of Student-t distributions for which random variable a is Gamma distributed; the proposed theorem extends thus their result to the class of all Gaussian scale mixtures for which $Ea^{-m} < \infty$.

5. APPLICATION: A RESULT BY FOLLAND

In this section, we show that the results above allow to recover easily the following theorem by Folland [5] about integration of a monomial over the sphere $\mathcal{S}_n$.

Theorem. *If*

$$P(X) = \prod_{i=1}^n X_i^{\alpha_i}$$

then

$$\int_{\mathcal{S}_n} P(X) d\sigma = \begin{cases} 0 & \text{if some } \alpha_i \text{ is odd} \\ \frac{2\Gamma(\beta_1)\dots\Gamma(\beta_n)}{\Gamma(\beta_1+\dots+\beta_n)} & \text{else} \end{cases}$$

where

$$\beta_i = \frac{\alpha_i + 1}{2}$$

and $d\sigma$ is the surface measure on the sphere $\mathcal{S}_n$.

Proof. Defining $\alpha_0 = 0$, we consider the function $T : [1, \sum_{i=1}^n \alpha_i] \rightarrow [1, n]$ such that

$$T(i) = j \quad \text{if} \quad \alpha_1 + \dots + \alpha_{j-1} < i \leq \alpha_1 + \dots + \alpha_j.$$

We choose forms $\mathcal{L}_i$ such that $l_i = \delta_{T(i)}$ where δ_k is the k -th column vector of the $n \times n$ identity matrix. Since the scalar product $l_i^t l_{\sigma(i)}$ equals zero when $T(i) \neq T(\sigma(i))$, we consider only the $\prod_{i=1}^n \frac{\alpha_i!}{2^{\frac{\alpha_i}{2}} (\frac{\alpha_i}{2})!}$ pairings for which $T(i) = T(\sigma(i))$ and $l_i^t l_{\sigma(i)} = \delta_{T(i)}^t \delta_{T(i)} = 1$. Thus, by Wick's theorem,

$$E \prod_{i=1}^n \mathcal{L}_i(U) = \frac{\Gamma(\frac{n}{2})}{2^m \Gamma(m + \frac{n}{2})} \prod_{i=1}^n \frac{\alpha_i!}{2^{\frac{\alpha_i}{2}} (\frac{\alpha_i}{2})!} = \frac{\Gamma(\frac{n}{2})}{2^m \Gamma(m + \frac{n}{2})} \frac{2^{\sum_{i=1}^n \beta_i}}{(2\pi)^{n/2}} \prod_{i=1}^n \Gamma(\beta_i).$$

Since $d\sigma$ is the unnormalized surface measure and as the surface of the sphere is $S(\mathcal{S}_n) = \frac{2\pi^{n/2}}{\Gamma(\frac{n}{2})}$,

$$\int_{\mathbb{R}^n} P(X) d\sigma = S(\mathcal{S}_n) \times E \prod_{i=1}^n \mathcal{L}_i(U)$$

which yields the result. $\square$

Acknowledgement. This work was performed during a visit by C. Vignat to S. Bhatnagar to the Computer Science and Automation Department of the Indian Institute of Sciences, Bangalore. C.V. thanks S. B. for his nice welcome.

REFERENCES

- [1] P. Repetowicz and P. Richmond, The Wick theorem for non-Gaussian distributions and its application for noise filtering of correlated q -Exponentially distributed random variables, unpublished, arXiv:math-ph/0411020 v1, Nov 2004
- [2] P. Etingof, Lecture Notes, Geometry and Quantum Field Theory, Fall 2002, MIT Open Courseware, <http://ocw.mit.edu/OcwWeb/Mathematics/18-238Fall2002/LectureNotes/index.htm>
- [3] K.-T. Fang, S. Kotz & K.W. Ng, Symmetric Multivariate and Related Distributions, Chapman and Hall, London, 1990
- [4] K.-C. Chu, Estimation and Decision for Linear Systems with Elliptical Random Processes, IEEE Tr. on Automatic Control, Oct 1973, pp. 500-505.
- [5] G. B. Folland, How to Integrate a Polynomial over a Sphere, The American Mathematical Monthly, Vol. 108-5, May 2001, pp. 446-448

INSTITUT GASPARD MONGE, UNIVERSITÉ DE MARNE LA VALLÉE, FRANCE AND DEPARTMENT OF COMPUTER SCIENCE AND AUTOMATION, INDIAN INSTITUTE OF SCIENCES, BANGALORE, INDIA

E-mail address: `vignat@univ-mlv.fr`, `shalabh@csa.iisc.ernet.in`